

		National Library of Medicine 		My NCBI [Sign In] [Regis]						
All Databases		PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Books		
Search PubMed		for				Go		Clear				
About Entrez Text Version Entrez PubMed Overview Help FAQ Tutorial New/Noteworthy E-Utilities PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut My NCBI (Cubby) Related Resources Order Documents NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central		<input checked="" type="checkbox"/> Limits		Preview/Index		History		Clipboard		Details		
		Limits: Publication Date to 1995/06/07										
		Display		Abstract		Show: 20		Sort		Send to Text		
		All: 1		Review: 0								
		<input type="checkbox"/> 1: Diabetes. 1994 Jan;43(1):33-9. Related Articles, Li										
<p>Isolation of nonobese diabetic mouse T-cells that recognize novel autoantigens involved in the early events of diabetes.</p> <p>Gelber C, Paborsky L, Singer S, McAteer D, Tisch R, Jolicœur C, Buelo R, McDevitt H, Fathman CG.</p> <p>ImmuLogic Pharmaceutical, Palo-Alto, CA 94304.</p> <p>Insulin-dependent diabetes mellitus (IDDM) is thought to result from chronic cell-mediated, autoimmune islet damage. Our aim was to identify the earliest T-cell autoantigen in IDDM, reasoning that this antigen could be causally involved in the initiation of the disease. Identification of the earliest beta-cell specific autoantigen is extremely important in allowing advances in prevention and treatment of initial events in the development of inflammatory insulinitis that precedes beta-cell destruction and overt diabetes. Therefore, we analyzed the proliferative responses of peripheral T-cells from young, female nonobese diabetic (NOD) mice to extracts of pancreatic beta-cell lines. We were able to demonstrate that T-cells responsive to beta-cell antigens exist in the peripheral lymphoid tissue of these mice in the absence of deliberate priming before the manifestation of histologically detectable insulinitis. T-cell lines and clones isolated from the peripheral lymphatic tissues of young, unimmunized, female NOD mice were also shown to react with extracts of beta-cells. Fractionation of the beta-cell extracts showed that these T-cell clones recognized multiple beta-cell-specific autoantigens but none of the previously reported putative autoantigens (glutamic acid decarboxylase [GAD]65, GAD67, Hsp65, insulin ICA 69, carboxypeptidase-H, and peripherin). Thus, we can conclude that the responses are specific for novel beta-cell autoantigens. Finally, NOD T-cell proliferative responses were also seen to an extract of human islets suggesting potential shared antigenic determinants between human and mouse beta-cells (ABSTRACT TRUNCATED AT 250 WORDS)</p> <p>PMID: 8262314 [PubMed - indexed for MEDLINE]</p>												